

Submillimetre spectroscopy of the volatile metabolome

F. Bondu¹

¹*Institut de Physique de Rennes, UMR 6251, CNRS/Université de Rennes 1*

Gaseous ensembles of molecules, called the volatilome, are fingerprints [1]–[3] of the metabolism of living organisms, including bacteria and archeobacteria. The submillimetre spectroscopy of these mixtures [4] could provide an interesting analysis alternative to gas chromatography coupled to mass spectroscopy (GC-MS) as there would be less effects (heating, ionization) damaging the integrity of large molecules. The characterization of ensembles would then in medical situations [5], [6] discriminate pathological situations or possibly identify bacteria resistant to antibiotics, and in food industry [7], [8] survey fermentation processes. The realization of an instrument meets several challenges. (a) The optical generation of submillimetre waves promises to attain a large frequency span, thus a large molecule span: we will present results towards such a compact and tuneable source. (b) The detection system should present a large signal to noise ratio, in order to detect molecules at trace levels. (c) The sampling system should take care that a large fraction of volatile biological molecules are actually liquid at ambient temperature and pressure.

References

- [1] A. Amann *et al.*, « The human volatilome: volatile organic compounds (VOCs) in exhaled breath, skin emanations, urine, feces and saliva », *J. Breath Res.*, vol. 8, n° 3, p. 034001, sept. 2014.
- [2] B. A. Kimball, « Volatile metabolome: problems and prospects », *Bioanalysis*, vol. 8, n° 19, p. 1987- 1991, oct. 2016.
- [3] M. Sohrabi, L. Zhang, K. Zhang, A. Ahmetagic, et M. Q. Wei, « Volatile Organic Compounds as Novel Markers for the Detection of Bacterial Infections », *Clin. Microbiol. Open Access*, vol. 03, n° 03, 2014.
- [4] A. M. Fosnight, B. L. Moran, D. R. Branco, J. R. Thomas, et I. R. Medvedev, « Chemical analysis of exhaled human breath using a novel sub-millimeter/terahertz spectroscopic approach », 2013, p. 1- 2.
- [5] J.-N. Cornu, G. Cancel-Tassin, V. Ondet, C. Girardet, et O. Cussenot, « Olfactory Detection of Prostate Cancer by Dogs Sniffing Urine: A Step Forward in Early Diagnosis », *Eur. Urol.*, vol. 59, n° 2, p. 197- 201, févr. 2011.
- [6] S. Bashiardes, H. Shapiro, S. Rozin, O. Shibolet, et E. Elinav, « Non-alcoholic fatty liver and the gut microbiota », *Mol. Metab.*, vol. 5, n° 9, p. 782- 794, sept. 2016.
- [7] A. Casaburi, P. Piombino, G.-J. Nychas, F. Villani, et D. Ercolini, « Bacterial populations and the volatilome associated to meat spoilage », *Food Microbiol.*, vol. 45, Part A, p. 83- 102, févr. 2015.
- [8] F. K. Tavoria, S. Dahl, F. J. Carballo, et F. X. Malcata, « Amino Acid Catabolism and Generation of Volatiles by Lactic Acid Bacteria », *J. Dairy Sci.*, vol. 85, n° 10, p. 2462- 2470, oct. 2002.